


# Discovery of the positron by Anderson, 1932

## Cosmic rays traversing a cloud chamber


Super-saturated water vapor  
Pb plate



# The Lamb Shift

$$\Delta\nu = 1058 \text{ MHz} = 0.035 \text{ cm}^{-1}$$

# Deviations from Dirac theory observed by Houston and Williams



Williams Phys.Rev. 54.558 (1938)

Dirac Theory

$$\nu = 0.364 \text{ cm}^{-1} \quad (f = 10.9 \text{ GHz})$$

$$\lambda = 2.74 \text{ cm (micro-wave)}$$

Note:  $1/\lambda = f[\text{GHz}]/29.97 [\text{cm}]$

from Williams Phys.Rev. 54.558 (1938)

$n=3$  to  $n=2$  is  $H\alpha$

$D\alpha$  deuterium  
line broadening is mostly  
doppler @100K

TABLE I. Recent measurements of the doublet interval in  $H\alpha$  and  $D\alpha$ .

INVESTIGATOR	YEAR	DOUBLET INTERVAL IN $\text{CM}^{-1}$	
		$H\alpha$	$D\alpha$
Houston and Hsieh <sup>1</sup>	1934	0.312	
Williams and Gibbs <sup>2</sup>	1934	0.304	0.317
Kopfermann <sup>3</sup>	1934		0.323
Spedding, Shane and Grace <sup>4</sup>	1934	0.314	0.318
Williams and Gibbs (Unpublished)	Dec. 1934	0.314	0.318
Heyden <sup>5</sup>	1937		0.331

<sup>1</sup> Houston and Hsieh, Phys. Rev. **45**, 263 (1934).

<sup>2</sup> Williams and Gibbs, Phys. Rev. **45**, 475 (1934).

<sup>3</sup> Kopfermann, Naturwiss. **22**, 218 (1934).

<sup>4</sup> Spedding, Shane and Grace, Phys. Rev. **47**, 38 (1935).

<sup>5</sup> Heyden, Zeits. f. Physik **106**, 499 (1937).



FIG. 3. Interferometer fringes of  $D\alpha$  obtained with a 5 mm étalon spacing.

$$5\text{mm} = \Delta\nu \text{ 1 cm}^{-1}$$




FIG. 4. Microphotometer tracing of part of the fringe pattern shown in Fig. 3.

Pasternak (1938) suggested that these results could be  $2S_{1/2}, 2P_{1/2}$  splitting of  $0.03 \text{ cm}^{-1}$ . However, most attributed discrepancies with Dirac to impurities in the source.

$$H_{\alpha} \text{ both } I(3D_{3/2}-2P_{1/2})/I(3P_{3/2}-2S_{1/2})=2.4$$

TABLE I.

Houston and Hsieh, Phys. Rev. 45, 263 (1934)

	H $\alpha$	H $\beta$	H $\gamma$	H $\delta$	H $\epsilon$
$\Delta\nu_{\text{theor}}$	0.319	0.344	0.353	0.358	0.360
$\Delta\nu_{\text{obs}}$	0.307	0.330	0.339	0.345	0.351
$\Delta\nu_{\text{corr}}$	0.308	0.330	0.339	0.343	0.345

3rd row:  $2S$  level displacement of  $0.030 \text{ cm}^{-1}$

# Houston and Hsieh, Phys. Rev. 45, 263 (1934)

TABLE I.

Line	$d_0$ (mm)	$\Delta\nu$ ( $\text{cm}^{-1}$ )
H $\beta$	7.58 $\pm$ 0.01	0.3298 $\pm$ 0.0004
H $\gamma$	7.38 $\pm$ .01	.3388 $\pm$ .0004
H $\delta$	7.245 $\pm$ .015	.3451 $\pm$ .0006
H $\epsilon$	7.13 $\pm$ .015	.3506 $\pm$ .0006

The method of measurement we have used is not directly applicable to H $\alpha$ , and so we have not given much attention to it. We have, however, made five measurements of three plates in the ordinary fashion, and have obtained  $\Delta\nu = 0.3171 \pm 0.0020$ . On the other hand, with an interferometer separation of 7.91 mm the minima were far from being equal. A rough estimate indicates that the minima would be equal at about 8.1 or 8.2 mm. If this could be applied directly to determine the separation of the centers of gravity of the lines, the separation would be 0.3086 or 0.3049. Of course this direct application is not justified in the case of H $\alpha$ , but the graphical analysis based on the theoretical form of this line, indicated that these results need to be increased by only about one percent to get the correct separation. Hence we must conclude that the method of direct visual

measurement gives results which are too large, at least for this line. This is not difficult to understand, since the peak of the line stands out much more on a plate than does the weaker companion, and so the measurement tends to be of the strong lines, rather than of the center of gravity. We hope to make a further analysis of H $\alpha$  immediately, but this observation explains the difference between our present results and previous work on these doublets.

## DISCUSSION OF THE RESULTS

Table II shows the position and the intensity of each of the components of the first five members of the Balmer series, as computed from the present theory, without the inclusion of the effect of a nuclear magnetic moment. The positions are those given by Sommerfeld's formula, which gives the energy levels permitted by Dirac's theory. The intensities are calculated by the methods of Sommerfeld and Unsöld.<sup>4, 6</sup> For small atomic numbers, this method gives the same results as a more rigorous calculation with relativistic functions.

<sup>6</sup> Saha and Banerji, Zeits. f. Physik **68**, 704 (1931), get the same results from Dirac's relativistic theory.



## Fine Structure of the Hydrogen Atom by a Microwave Method\* \*\*

WILLIS E. LAMB, JR. AND ROBERT C. RETHERFORD

*Columbia Radiation Laboratory, Department of Physics, Columbia University, New York, New York*

(Received June 18, 1947)

THE spectrum of the simplest atom, hydrogen, has a fine structure<sup>1</sup> which according to the Dirac wave equation for an electron moving in a Coulomb field is due to the combined effects of relativistic variation of mass with velocity and spin-orbit coupling. It has been considered one of the great triumphs of Dirac's theory that it gave the "right" fine structure of the energy levels. However, the experimental attempts to obtain a really detailed confirmation through a study of the Balmer lines have been frustrated by the large Doppler effect of the lines in comparison to the small splitting of the lower or  $n = 2$  states. The various spectroscopic workers have alternated between finding confirmation<sup>2</sup> of the theory and discrepancies<sup>3</sup> of as much as eight percent. More accurate information would clearly provide a delicate test of the form of the correct relativistic wave equation, as well as information on the possibility of line shifts due to coupling of the atom with the radiation field and clues to the nature of any non-Coulombic interaction between the elementary particles: electron and proton.


# radar technology

During World War II, Willis worked on radar at Columbia. That expertise, together with his deep knowledge of quantum theory, put him in a good position to carry out his famous level-shift (that is, the Lamb shift) measurements in the hydrogen atom shortly after the war.

February 1940, Great Britain developed the [resonant-cavity magnetron](#), capable of producing microwave power in the kilowatt range, opening the path to second-generation radar systems.<sup>[4]</sup>.... Bell Labs was able to duplicate the performance, and the [Radiation Laboratory](#) at MIT was established to develop microwave radars. (Wikipedia)



# Lamb Nobel 1955



Electric dipole atomic transition rules:


$$\Delta l = 1, -1 \quad \Delta m_l = 1, 0, -1 \quad \Delta s = 0 \quad \text{allowed } \tau \sim \text{ns}$$

$$\Delta j = 1, 0, -1 \quad \text{no } j=0 \text{ to } j=0 \quad 2S_{1/2} \leftrightarrow 1S_{1/2} \quad \tau = 1/7 \text{ s}$$

Micro-wave induced transitions:

$$\alpha \text{ to } a, b, c \quad \beta \text{ to } b, c, d \quad \alpha, \beta \text{ to } e \quad \alpha, \beta \text{ to } f$$

# schematic of apparatus



The excited atoms passed through a region containing both microwave radiation and an adjustable magnetic field, and then hit a metal foil. The excited atoms ( $2S_{1/2}$ ) would, upon hitting the foil, drop back to the ground state emitting electrons that the team could detect as a current. The key to the experiment was that if the magnetic-field-induced energy difference between the two states was equal to the energy of the microwave photons, then the long-lived  $2S_{1/2}$  state would absorb a photon and turn into the short-lived  $P$ -state. These atoms would drop back to the  $1S_{1/2}$  ground state before reaching the target, and the current in the detector would essentially vanish.




Fig. 3. Cross section of second apparatus: (a) tungsten oven of hydrogen dissociator, (b) movable slits, (c) electron bombarder cathode, (d) grid, (e) anode, (f) transmission line, (g) slots for passage of metastable atoms through interaction space, (h) plate attached to center conductor of r-f transmission line, (i) d.c. quenching electrode, (j) target for metastable atoms, (k) collector for electrons ejected from target, (l) pole face of magnet, (m) window for observation of tungsten oven temperature.




FIG. 1. A typical plot of galvanometer deflection due to interruption of the microwave radiation as a function of magnetic field. The magnetic field was calibrated with a flip coil and may be subject to some error which can be largely eliminated in a more refined apparatus. The width of the curves is probably due to the following causes: (1) the radiative line width of about 100 Mc/sec. of the  $^2P$  states, (2) hyperfine splitting of the  $^2S$  state which amounts to about 88 Mc/sec., (3) the use of an excessive intensity of radiation which gives increased absorption in the wings of the lines, and (4) inhomogeneity of the magnetic field. No transitions from the state  $2^2S_{\frac{1}{2}}(m = -\frac{1}{2})$  have been observed, but atoms in this state may be quenched by stray electric fields because of the more nearly exact degeneracy with the Zeeman pattern of the  $^2P$  states.

Lamb, 1947

observed  
current  $\sim 10^{-14}$  A



Lamb, 1947





FIG. 2. Experimental values for resonance magnetic fields for various frequencies are shown by circles. The solid curves show three of the theoretically expected variations, and the broken curves are obtained by shifting these down by 1000 Mc/sec. This is done merely for the sake of comparison, and it is not implied that this would represent a "best fit." The plot covers only a small range of the frequency and magnetic field scale covered by our data, but a complete plot would not show up clearly on a small scale, and the shift indicated by the remainder of the data is quite compatible with a shift of 1000 Mc.

$2S_{1/2} \rightarrow 2P_{3/2}$  shifted down from expected if  $2S_{1/2}, 2P_{1/2}$  were degenerate, implying  $E(2S_{1/2}) > E(2P_{1/2})$  by 1000MHz






Observed lines  
Versus magnetic field

Lamb Nobel 1955

M. Gold

physics 522, Spring 2025

# Lamb Nobel 1955



## The Electromagnetic Shift of Energy Levels

H. A. BETHE

*Cornell University, Ithaca, New York*

(Received June 27, 1947)

**B**Y very beautiful experiments, Lamb and Retherford<sup>1</sup> have shown that the fine structure of the second quantum state of hydrogen does not agree with the prediction of the Dirac theory. The  $2s$  level, which according to Dirac's theory should coincide with the  $2p_{1/2}$  level, is actually higher than the latter by an amount of about  $0.033 \text{ cm}^{-1}$  or 1000 megacycles. This discrepancy had long been suspected from spectroscopic measurements.<sup>2,3</sup> However, so far no satisfactory theoretical explanation has been given. Kemble and Present, and Pasternack<sup>4</sup> have shown that the shift of the  $2s$  level cannot be

explained by a nuclear interaction of reasonable magnitude, and Uehling<sup>5</sup> has investigated the effect of the "polarization of the vacuum" in the Dirac hole theory, and has found that this effect also is much too small and has, in addition, the wrong sign.

Schwinger and Weisskopf, and Oppenheimer have suggested that a possible explanation might be the shift of energy levels by the interaction of the electron with the radiation field. This shift comes out infinite in all existing theories, and has therefore always been ignored. However, it is possible to identify the most strongly (linearly) divergent term in the level shift with an electromagnetic *mass* effect which must exist for a bound as well as for a free electron. This effect should

<sup>1</sup> Phys. Rev. **72**, 241 (1947).

<sup>2</sup> W. V. Houston, Phys. Rev. **51**, 446 (1937).

<sup>3</sup> R. C. Williams, Phys. Rev. **54**, 558 (1938).

<sup>4</sup> E. C. Kemble and R. D. Present, Phys. Rev. **44**, 1031 (1932); S. Pasternack, Phys. Rev. **54**, 1113 (1938).

<sup>5</sup> E. A. Uehling, Phys. Rev. **48**, 55 (1935).